

Fertilizer Contains Dimension* 0.1% FG

Turf and Ornamental Herbicide

For season-long control of crabgrass and control or suppression of listed annual grasses and broadleaf weeds in ornamental landscape areas, established lawns and ornamental turf, including golf course fairways, roughs and tee boxes.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

In the state of New York, this product may be applied only by commercial applicators at no more than 500 lb (0.5 lb active ingredient) per acre (11.5 lb per 1000 sq ft) per year. Use of this product in Nassau and Suffolk counties of New York is prohibited.

Active Ingredients: dithiopyr: 3,5-pyridinedicarbothioic acid,2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S, S-dimethyl ester00.1%
Inert Ingredients.....99.9%
Total.....100.0%

19-0-6

Guaranteed Fertilizer Analysis

Total Nitrogen (N) 19.00%
 19.00% Urea Nitrogen*
 Soluble Potash (K₂O) 6.00%
 Sulfur (S) 2.62%
 2.62% Elemental Sulfur (S)

Derived from: Urea, Muriate of Potash, and Poly Coated Sulfur Coated Urea.
 *5.7% Slowly Available Urea Nitrogen from Sulfur Coated Urea.

E.P.A. REG. NO. 62719-486-41124

E.P.A. EST. No. 41124-WI-1 (WI)
 41124-OH-1 (OH)

The letter of production batch code indicates producing establishment
ITEM NO. 6000437 **NET WT. 50 LB**

*Trademark of Dow AgroSciences LLC
 Dow AgroSciences LLC • Indianapolis, Indiana 46268 USA

Table 1 Recommended Use Rates

Region	Application Rates	Program 1	Program 2	Program 3
All states, except NY, and parts of states not included in transition, south, coastal south or west.	lb/acre	190	250	380
	lb/1000 sq ft	4.4	5.7	8.7
	lb ai/acre	0.19	0.25	0.38
Transition: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, & RI	lb/acre	250	380	500
	lb/1000 sq ft	5.7	8.7	11.5
	lb ai/acre	0.25	0.38	0.5
South: AL, AR, GA, LA, MS, NC, NM, OK, SC, TN, & TX	lb/acre	380	250 + 250	310 + 310
	lb/1000 sq ft	8.7	5.7 + 5.7 [†]	7.1 + 7.1 [†]
	lb ai/acre	0.38	0.25 + 0.25	0.31 + 0.31
Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX	lb/acre	250 + 250	310 + 310	380 + 380
	lb/1000 sq ft	5.7 + 5.7 [†]	7.1 + 7.1 [†]	8.7 + 8.7 [†]
	lb ai/acre	0.25 + 0.25	0.31 + 0.31	0.38 + 0.38
West: AZ, CA, & NV. In this climatically diverse region, use the higher rates in local areas with longer crabgrass seasons.	lb/acre	190 + 250	250 + 380	250 + 250
	lb/1000 sq ft	4.4 + 5.7	5.7 + 8.7	5.7 + 5.7 [†]
	lb ai/acre	0.19 + 0.25	0.25 + 0.38	0.25 + 0.25

[†]Split applications may be made 5 - 10 weeks apart.