

FERTILIZER

With



AND



Active Ingredients:

isoxaben: <i>N</i> -[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomers...	0.38%
trifluralin: <i>a,a</i> -a-trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine.....	0.39%
benefin: <i>N</i> -butyl- <i>N</i> -ethyl- <i>a,a,a</i> -trifluoro-2,6-dinitro- <i>p</i> -toluidine.....	0.76%
Inert Ingredients.....	98.47%
Total.....	100.00%

Contains 0.76 lb active ingredient per 50 lb bag.

This fertilizer product contains Gallery and Team preemergence herbicides for control of certain broadleaf weeds and annual grasses in established turfgrasses and landscape ornamentals. This product will not control established weeds. Existing weeds should be controlled by cultivation or with postemergence herbicides.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994

Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.

©™Trademark of Dow AgroSciences LLC

Dow AgroSciences LLC • Indianapolis, Indiana 46268 U.S.A.

Keep Out of Reach of Children

CAUTION

Hazards to Humans and Domestic Animals

Precautionary Statements

Harmful if Swallowed or Inhaled • **Prolonged or Frequently Repeated Skin Contact May Cause Allergic Reactions in Some Individuals**

Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment:

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates, including shrimp and oyster. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Read all directions carefully before applying this product.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow others to enter the treated areas until dusts have settled. If soil incorporation is required after the application, do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. If the incorporation is accomplished by watering-in, do not enter or allow others to enter the treated area until the surface is dry after the watering-in.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior landscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

5-15-15

GUARANTEED ANALYSIS

Total Nitrogen (N).....	5.0%
3.2% Ammoniacal Nitrogen	
1.8% Water Insoluble Nitrogen	
Available Phosphate (P2O5).....	15.0%
Soluble Potash (K2O).....	15.0%
Sulfur (S)	5.0%
Iron (Fe)	1.0%

Derived From: BioSolids, Ammoniated Phosphate, and Sulfate of Potash.

General Information

This fertilizer product contains Gallery® and Team™, preemergence herbicides for control of certain broadleaf weeds and annual grasses in established turfgrass and landscape ornamentals. This product will not control existing weeds. Existing weeds should be controlled by cultivation or with a foliar applied postemergence herbicide.

Optimum weed control performance will be obtained when an application is followed by overhead irrigation or rainfall within a few hours. A single rainfall or sprinkler irrigation of 0.5 inch or more is required to activate this product.

Use Precautions and Restrictions:

- Apply to newly seeded turf (including overseeded turf) only after seedlings are established (three leaf stage and tillering) and well rooted. Do not overseed established turf sooner than 60 days following an application.
- **Reseeding:** Do not reseed treated areas for a minimum of 8 weeks after a single application of 175 lb per acre, or 12 to 16 weeks after a single application of 265 lb per acre.
- Do not apply to areas where desirable plants will be grown from seed. Use only on established perennial plants. Take precautions to prevent inadvertent application from rotary-type applicators to such areas. Herbicides in this product prevent seed emergence and may injure bedding plants.
- This product may injure turf that is weakened due to severe winter weather or is not well established.
- Do not apply to newly sprigged or hydromulched areas of Bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until turf is well established.
- Do not apply to dichondra.
- Do not apply to golf course putting greens.
- Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of this product.
- Do not overdose or overlap during application in such a way as to apply in excess of labeled rates as turf injury may occur.

Treatment of Turf or Ornamental Species Not Listed on the Label for T&O Fertilizer – Contains Gallery Plus Team

Users who wish to use this product on species not listed on this label may determine the suitability for such uses by treating a small area or a small number of plants at a recommended rate. Prior to treating a larger area, the treated area/plants should be observed for any sign of herbicidal injury during 30 to 60 days of typical growing conditions. **The user assumes responsibility for any plant damage or other liability resulting from use of this product on species not recommended on this label.**

This product is recommended for use on the following established perennial turfgrass species:

Established Cool Season Grasses

Common Name	Scientific Name
bentgrass, colonial	<i>Agrostis tenuis</i>
bentgrass, creeping	<i>Agrostis stolonifera</i>
bluegrass, Kentucky	<i>Poa pratensis</i>
fescue, chewings	<i>Festuca rubra</i> var. <i>commutata</i>
fescue, creeping red	<i>Festuca rubra</i>
fescue, sheep	<i>Festuca ovina</i>
fescue, tall	<i>Festuca arundinaceae</i>
ryegrass, perennial	<i>Lolium perenne</i>

Established Warm Season Grasses

Common Name	Scientific Name
bahiagrass	<i>Paspalum notatum</i> var. <i>Saurae parodi</i>
Bermudagrass	<i>Cynodon dactylon</i>
centipedegrass	<i>Eremochloa ophiuroides</i>
fescue, tall	<i>Festuca arundinaceae</i>
St. Ausustinegrass	<i>Stenotaphrum secundatum</i>
zoysiagrass	<i>Zoysia japonica</i>
zoysiagrass	<i>Zoysia tenuifolia</i>

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container. Protect stored product from moisture. In case of leak or spill, contain material and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

E.P.A. REG. NO. 62719-280-41124

E.P.A. EST. NO. 41124-WI-1

NET WT. 50 LBS. / COVERS 8,333 SQ. FT.

This product may be used for annual weed control in landscape settings on the following established ornamental plant species:

Ornamental Trees

Common Name	Scientific Name
almond	<i>Prunus dulcis</i>
apple, crabapple	<i>Malus</i> spp.
apricot	<i>Prunus armeniaca</i>
birch, paper	<i>Betula papyrifera</i>
cherry	<i>Prunus</i> spp.
cottonwood	<i>Populus deltoides</i>
dogwood, flowering	<i>Cornus florida</i>
dogwood, kousa	<i>Cornus kousa</i>
honeylocust	<i>Gleditsia triacanthos</i>
maple, flowering	<i>Abutilon hybridum</i>
maple, red	<i>Acer rubrum</i>
maple, silver	<i>Acer saccharinum</i>
oak, pin	<i>Quercus palustris</i>
peach	<i>Prunus persica</i>
pine, Austrian	<i>Pinus nigra</i>
pine, eastern white	<i>Pinus strobus</i>
pine, Japanese black	<i>Pinus thunbergiana</i>
pine, scotch	<i>Pinus sylvestris</i>
plum	<i>Prunus</i> spp.
redbud, eastern	<i>Cercis canadensis</i>
spruce, Colorado	<i>Picea pungens</i>
spruce, Norway	<i>Picea abies</i>
sweetgum	<i>Liquidambar styraciflua</i>
sycamore	<i>Platanus occidentalis</i>
walnut, black	<i>Juglans nigra</i>

Ornamental Shrubs

Common Name	Scientific Name
arborvitae	<i>Thuja occidentalis</i>
azalea	<i>Rhododendron</i> spp.
boxwood, littleleaf	<i>Buxus microphylla</i>
camellia, Japanese	<i>Camellia japonica</i>
camellia, sasanqua	<i>Camellia sasanqua</i>
cinquefoil	<i>Potentilla fruticosa</i>
cleysera, Japanese	<i>Cleyera japonica</i>
cotoneaster, cranberry	<i>Cotoneaster apiculata</i>
elaeanus, silverberry	<i>Elaeagnus pungens</i>
euonymus, winged	<i>Euonymus alata</i>
euonymus, wintercreeper	<i>Euonymus fortunei</i>
forsythia	<i>Forsythia</i> spp.
holly	<i>Ilex</i> spp.
honeysuckle	<i>Lonicera</i> spp.
juniper, creeping	<i>Juniperus horizontalis</i>
laurel, mountain	<i>Kalmia latifolia</i>
lilac, common	<i>Syringa vulgaris</i>
myrtle, wax	<i>Myrica cerifera</i>
privet	<i>Ligustrum</i> spp.
pyracantha	<i>Pyracantha fortuneana</i>
redcedar, eastern	<i>Juniperus virginiana</i>
rhododendron	<i>Rhododendron</i> spp.
spiraea, vanhouttei	<i>Spiraea vanhouttei</i>
viburnum	<i>Viburnum</i> spp.
yew, Japanese	<i>Taxus cuspidata</i>
yewpine	<i>Podocarpus macrophyllum</i>

Ornamental Ground Cover Plantings

Common Name	Scientific Name
ceanothus	<i>Ceanothus</i> spp.
coyote brush	<i>Baccharis pilularis</i>
daisy trailing African	<i>Osteospermum fruticosum</i>
gazania	<i>Gazania</i> spp.
ice plant, largeleaf	<i>Carpobrotus edulis</i>
ivy, Algerian	<i>Hedera canariensis</i>
ivy, English	<i>Hedera helix</i>
lilyturf, bigblue	<i>Liriope muscari</i>
plumbago, dwarf	<i>Ceratostigma umbaginoides</i>
rosemary	<i>Rosmarinus officinalis</i>
snow-in-summer	<i>Cerastium tomentosum</i>
St. Johnswort	<i>Hypericum coris</i>

Application Directions

Application Equipment and Application Techniques

Apply this product using a drop or rotary-type spreader designed to apply granular fertilizer. Avoid the use of spreaders that apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate. Apply this product uniformly over the lawn. A more uniform application may be achieved by spreading half of the required amount over the area and then applying the remaining half at right angles to the previous direction. Avoid skips or overlaps as poor weed control or turf injury may occur. Check frequently to be sure equipment is operating properly and distributing product uniformly. A splash board mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Broadcast Application Rates:

Season of Application	Pounds per Acre	Pounds per 1000 sq ft
Spring Application (Summer annuals)	175 to 265 lb/acre	4.0 to 6.0 lb/1000 sq ft
Fall Application (Winter annuals)	265 lb/acre	6.0 lb/1000 sq ft

Item No. 2003800

052305

Distributed By:

Vogel Seed & Fertilizer, Inc. • Jackson, Wisconsin, 53037

Weeds Controlled

Weeds controlled when applied prior to weed germination:

Common Name	Scientific Name
annual bluegrass	<i>Poa annua</i>
barnyardgrass	<i>Echinochloa crus-galli</i>
burweed, lawn	<i>Soliva pterosperma</i>
buttercup	<i>Ranunculus</i> spp.
carpetweed	<i>Mollugo verticillata</i>
celery, wild	<i>Apium leptophyllum</i>
chickweed, common	<i>Stellaria media</i>
chickweed, mouseear	<i>Cerastium vulgatum</i>
crabgrass	<i>Digitaria</i> spp.
(large crabgrass)	
(smooth crabgrass)	
foolsparsley	<i>Aethusa cynapium</i>
foxtails	<i>Setaria</i> spp.
(bottlegrass)	
(bristlegrass)	
(giant foxtail)	
(green foxtail)	
(pigeongrass)	
(robust foxtail)	
geranium, Carolina	<i>Geranium carolinianum</i>
goosegrass	<i>Eleusine indica</i>
henbit <i>Lamium amplexicaule</i>	
knotted, prostrate	<i>Polygonum aviculare</i>
mallow, little	<i>Malva parviflora</i>
parsley - piert	<i>Alchemilla arvensis</i>
pennywort, lawn	<i>Hydrocotyle sibthorpioides</i>
pineappleweed	<i>Matricaria matricarioides</i>
speedwell, thymeleaf	<i>Veronica serpyllifolia</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
spurge, spotted	<i>Euphorbia maculata</i>
spurry, corn	<i>Spergula arvensis</i>
woodsorrel, creeping	<i>Oxalis corniculata</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

Weeds suppressed or partially controlled by a single application or controlled by repeat applications when applied prior to weed germination:

Common Name	Scientific Name
clover spp.	<i>Trifolium</i> spp.
clover, white ¹	<i>Trifolium repens</i>
dandelion ¹	<i>Taraxacum officinale</i>
groundsel, common	<i>Senecio vulgaris</i>
medic, black	<i>Medicago lupulina</i>
plantain, broadleaf	<i>Plantago major</i>
plantain, buckhorn	<i>Plantago lanceolata</i>
plantain, slender	<i>Plantago elongata</i>

¹From seed only

Uses

Established Turfgrass

Application Timing: Apply in late summer or early fall approximately 3 weeks prior to germination of winter annual weeds, or in early spring 3 weeks prior to germination of weeds which emerge in spring or early summer.

Broadcast Application Rates

Apply this product at a rate of 175 to 265 lb per acre (4 to 6 lb per 1000 sq ft). A minimum of 175 lb per acre (4 lb per 1000 sq ft) is recommended for optimum results. Use the higher rate in the rate range where heavy weed infestation is anticipated.

Established Ornamental Plants in Landscape Settings

This product is recommended as a preemergence application for control of certain broadleaf weeds and annual grasses in a wide variety of established ornamental trees, shrubs, and ground covers. Apply only to non-bearing fruit and nut trees and non-bearing vineyards which are defined as plants that will not bear fruit for at least 1 year after treatment.

Application Timing: Apply 1 to 2 weeks prior to the germination of weed seeds or clean-cultivate and hoe to remove existing weeds before applying this product. Apply to the smooth soil surface (no large clods) to ensure uniform distribution around established plants or on ornamental beds. Do not apply this product to plantings when the foliage is wet.

Broadcast Application Rates

Apply this product at a rate of 175 to 265 lb per acre (4 to 6 lb per 1000 sq ft) as a spring application or 265 lb per acre (6 lb per 1000 sq ft) as a fall application. In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 to 12 weeks following the initial application. Use the higher rate in the rate range where heavy weed infestation is anticipated.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.