

FERTILIZER

Contains  Pro

A granular fertilizer containing Team Pro herbicide for the preemergence control of crabgrass, foxtail, goosegrass, Poa annua and other annual grass and broadleaf weeds in established turfgrass.

Active Ingredients:
 benefin: *N*-butyl-*N*-ethyl-*a,a,a*-trifluoro-2,6-dinitro-*p*-toluidine..... 0.43%
 trifluralin: *a,a,a*-trifluoro-2,6-dinitro-*N,N*-dipropyl-*p*-toluidine..... 0.43%
 Inert Ingredients:.....99.14%
 Total.....100.00%

Keep Out of Reach of Children

CAUTION

Notice: Read the entire label. Use only according to label directions.

In case of emergency endangering health or the environment involving this product, call collect 517-636-4400.

Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.

E.P.A. REG. NO. 62719-289-41124 E.P.A. EST. NO. 41124-WI-1, EST NO. 41124-OH-1

First letter of lot code indicates production establishment

* Trademark of Dow Agrosciences

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation. Harmful If Swallowed, Inhaled Or Absorbed Through The Skin
 Avoid breathing dust and contact with eyes, skin or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use protective clothing such as coveralls, a long-sleeved shirt and long pants, shoes and socks, and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

FIRST AID

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.
If swallowed: Do not induce vomiting. Call physician or Poison Control Center. Drink promptly a large quantity of milk, egg white, gelatin solution, or if these are not available, large quantities of water. Avoid alcohol.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation develops.

If inhaled: If breathing difficulty occurs, move individual to fresh air and get medical attention. If not breathing, provide cardiopulmonary resuscitation assistance (mouth to mouth) and get medical attention immediately.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

20-0-8

GUARANTEED ANALYSIS

Total Nitrogen (N).....	20.00%
18.50% Urea Nitrogen*	
0.15% Ammoniacal Nitrogen	
1.35% Water Insoluble Nitrogen	
Soluble Potash (K ₂ O).....	8.00%
Sulfur(S).....	0.37%
Iron (Fe).....	1.00%
Derived From: Bio-Solids, Ammoniated Phosphate, Muriate of Potash, Urea, and Polymer- Coated, Sulfur- Coated Urea.	
*3.72% slowly available water-soluble nitrogen derived from Polymer-Coated, Sulfur-Coated Urea.	

Weeds Controlled

Annual Grasses:

Common Name
 annual bluegrass
 crabgrass, smooth
 crabgrass, large or hairy
 goosegrass (silver crabgrass or crowfootgrass)
 barnyardgrass (watergrass)
 green foxtail
 yellow foxtail

Broadleaf Weeds:

Common Name
 chickweed, common
 knotweed, prostrate
 lambsquarters, common
 pigweed
 purslane, common

Scientific Name

Poa annua
Digitaria ischaemum
Digitaria sanguinalis
Eleusine indica
Echinochloa crus-galli
Setaria viridis
Setaria glauca

Scientific Name

Stellaria media
Polygonum aviculare
Chenopodium album
Amaranthus spp.
Portulaca oleracea

Weeds Suppressed:

Common Name

panicum, Texas
 pusley, Florida
 prostrate spurge
 spotted spurge
 creeping spurge
 creeping woodsorrel
 woodsorrel

Scientific Name

Panicum texanum
Richardia scabra
Euphorbia humistrata
Euphorbia maculata
Euphorbia serpens
Oxalis corniculata
Oxalis spp.

Approved Uses

Established Turfgrass

When to Apply: For optimum results, this product must be applied one to two weeks prior to the anticipated germination of the target annual weeds. A single application will usually provide season-long weed control. However, a second or split application is recommended when the initial application is made well in advance of weed germination or when extended periods of weed control is required, such as areas with longer growing seasons or for lawn care operator programs.

How to Apply: Apply this product evenly over the turfgrass area using a properly calibrated drop type or rotary spreader. Calibrate the spreader prior to use according to the manufacturer's directions. Check frequently to make sure equipment is working properly and distributing granules uniformly. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. Avoid use of spreaders that apply material in narrow concentrated bands. A splashboard mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Activation: Optimum weed control performance will be obtained if treated areas receive rainfall or irrigation soon after application.

Important Use Precautions: Do not apply this product to golf course putting greens, dichondra, or newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass. Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application. This product will not harm most nearby ornamentals.

This Product may be applied to the following established turfgrass species:

Established Cool Season Turfgrasses	Established Warm Season Turfgrasses
Bentgrass fairways	Bahiagrass
Kentucky bluegrass	Bermudagrass
Perennial ryegrass	Centipedegrass
Fescues	Fescue, tall
	St. Augustinegrass
	Zoysiagrass

Broadcast Application Rates:

Application Timing
Spring: To control spring-to-summer germinating annual weeds.
Fall: To control fall-germinating weeds

Application Rate
174 to 349 lb/acre
(4.00 to 8.01 lb/1000 sq ft)

Specific Use Directions

In cool season and warm season turfgrass areas, make a single application in the spring or fall, 1 to 2 weeks before expected time of weed germination.

Split Application:

For extended control of spring and summer annual weeds

174 lb/acre followed by **174 lb/acre**
(4.00 followed by 4.00 lb/1000 sq ft)

In Cool season turfgrass area and northern half of the warm season turfgrass area, apply the first application in spring 1 to 2 weeks before expected time of weed germination and make a second application 8 to 10 weeks later.

233 lb/acre followed by **233 lb/acre**
(5.34 followed by 5.34 lb/1000 sq ft)

In southern half of warm season turfgrass area, apply the first application in spring 1 to 2 weeks before the expected time of weed germination and make a second application 10 to 12 weeks later.

Reseeding Precautions

Team herbicide, which controls annual grass weeds, may also affect the establishment of new desirable turfgrass seedlings, if reseeding is attempted too soon after application. When reseeding, it is essential that proper cultural practices, such as soil cultivation, irrigation and fertilization be followed. Reseeding, following the last application of this product, should not be attempted until the applicable minimum time delay in the following table has been met:

Amount of Product Applied Per Acre (Pounds per 1000 sq ft)	Minimum Time Delay Following Last Application Before Reseeding (Weeks)
Single application of 174 lb/acre (4.00 lb/1000 sq ft)	8
Single application of 349 lb/acre (8.01 lb/1000 sq ft)	12 to 16
Split application of 174 lb/acre (4.00 lb/1000 sq ft)	8 to 10
Split application of 233 lb/acre (5.34 lb/1000 sq ft)	10 to 12

Established Ornamental Plants in Landscape Settings

This product is recommended for preemergence control of annual grasses and certain broadleaf weeds in a wide variety of established ornamental trees, shrubs, and flowers.

Application timing: Apply 1 to 2 weeks prior to the expected germination time of target annual weeds or to clean cultivated weed free areas. Do not apply this product to plantings when the foliage is wet.

Soil Preparation: This product can be surface applied around established plants or on ornamental beds. Apply to a smooth soil surface (no large clods) to assure uniform distribution. Clean-cultivate ad hoe to remove existing weeds before applying this product.

Activation: Optimum weed control performance will be obtained when followed by overhead irrigation or rainfall, within a few hours after application.

Broadcast Application Rates:

Season of Application	Pounds per 1000 sq ft
Spring Application (Summer annuals)	4.00 to 8.01 lb/1000 sq ft
Fall Application (Winter annuals)	4.00 to 8.01 lb/1000 sq ft

In areas with extended growing seasons or where desired, a second application at the recommended rate can be made approximately 10 - 12 weeks following the initial application.

NET WT. 50 LBS.

COVERS 12,500 SQ. FT.

Distributed By:
 Vogel Seed & Fertilizer, Inc.
 Jackson, Wisconsin, 53037

Item No. 2247740

042408re